

# Molecular fraction calculations for an atomic-molecular Bose-Einstein condensate model

Jon Links

Centre for Mathematical Physics,  
The University of Queensland,  
Australia.

2nd Annual Meeting of ANZAMP,  
Mooloolaba 2013

# Hamiltonian

The model consists of three interacting bosonic degrees of freedom:

$$H = U_{aa}N_a^2 + U_{bb}N_b^2 + U_{cc}N_c^2 + U_{ab}N_aN_b + U_{ac}N_aN_c + U_{bc}N_bN_c \\ + \mu_aN_a + \mu_bN_b + \mu_cN_c + \Omega(a^\dagger b^\dagger c + c^\dagger ba).$$

It commutes with the total atom number  $N = N_a + N_b + 2N_c$  and the atomic imbalance  $J = N_a - N_b$ . We introduce

$$k = J/N, k \in [-1, 1]$$

as the fractional atomic imbalance. The classical analogue is the non-linear pendulum

$$H = \lambda p_\phi^2 + 2(\alpha - \lambda)p_\phi + \lambda - 2\alpha + \beta \\ + \sqrt{2(1 - p_\phi)(p_\phi + c_+)(p_\phi + c_-)} \cos\left(\frac{4\phi}{N}\right)$$

with

$$\lambda = \frac{\sqrt{2N}}{\Omega} \left( \frac{U_{aa}}{4} + \frac{U_{bb}}{4} + \frac{U_{cc}}{4} + \frac{U_{ab}}{4} - \frac{U_{ac}}{4} - \frac{U_{bc}}{4} \right)$$

$$\alpha = \frac{\sqrt{2N}}{\Omega} \left( \frac{1+k}{2} U_{aa} + \frac{1-k}{2} U_{bb} + \frac{1}{2} U_{ab} - \frac{1+k}{4} U_{ac} - \frac{1-k}{4} U_{bc} \right. \\ \left. + \frac{1}{2N} (\mu_a + \mu_b - \mu_c) \right)$$

$$\beta = \frac{\sqrt{2N}}{\Omega} \left( (1+k)^2 U_{aa} + (1-k)^2 U_{bb} + (1-k^2) U_{ab} \right. \\ \left. + \frac{2}{N} ((1+k)\mu_a + (1-k)\mu_b) \right)$$

$$c_{\pm} = 1 \pm 2k.$$

# Order parameter - the molecular fraction

Define the order parameter

$$\mathcal{O} = 2 \frac{\langle N_c \rangle}{N} = \frac{2}{N} \frac{\partial E_0}{\partial \mu_c}$$

which measures the average molecular fraction:

$$\begin{array}{ll} \mathcal{O} = 0 & \text{atomic phase,} \\ 0 < \mathcal{O} < 1 & \text{mixed phase,} \\ \mathcal{O} = 1 & \text{molecular phase.} \end{array}$$

The order parameter relates to the momentum of the classical system through

$$\mathcal{O} \mapsto \frac{1}{2} (1 - p_\phi).$$

It is not associated with symmetry breaking.

# Classical fixed points

The dynamical evolution is governed by Hamilton's equations:

$$\begin{aligned} \frac{dp_\phi}{dt} &= \frac{\partial H}{\partial \phi} = -\frac{4}{N} \sqrt{2(1-p_\phi)(p_\phi+c_+)(p_\phi+c_-)} \sin\left(\frac{4\phi}{N}\right), \\ -\frac{d\phi}{dt} &= \frac{\partial H}{\partial p_\phi} = 2\lambda p_\phi + 2\alpha - 2\lambda \\ &\quad + \frac{(1-p_\phi)(2p_\phi+2) - (p_\phi+c_+)(p_\phi+c_-)}{\sqrt{2(1-p_\phi)(p_\phi+c_+)(p_\phi+c_-)}} \cos\left(\frac{4\phi}{N}\right). \end{aligned}$$

The fixed points of the system are determined by the condition

$$\frac{\partial H}{\partial \phi} = \frac{\partial H}{\partial p_\phi} = 0.$$

Phase boundaries of the parameter space are identified according to fixed point bifurcations.






Molecular fraction:  $\lambda = \mu_a = \mu_b = 0$  - numerical

Figure : Order parameter. Inset shows the first derivative.



# Bethe ansatz solution

$$E = AM(M - 1) + BM + C - \Omega \sum_{j=1}^M v_j,$$

where for  $J \geq 0$

$$\frac{\Omega(J + 1 - v_j^2) + Bv_j}{v_j(\Omega + Av_j)} = \sum_{k \neq j}^M \frac{2}{v_k - v_j}, \quad j = 1, 2, \dots, M,$$

$M = (N - J)/2$ ,  $L = (N + J)/2$ , and

$$A = U_{aa} + U_{bb} + U_{cc} + U_{ab} - U_{ac} - U_{bc},$$

$$B = (1 + 2L - 2M)U_{aa} + U_{bb} + (1 - 2M)U_{cc} + (1 + L - M)U_{ab} \\ + (2M - L - 1)U_{ac} + (M - 1)U_{bc} + \mu_a + \mu_b - \mu_c,$$

$$C = (L - M)^2 U_{aa} + M(L - M)U_{ac} + M^2 U_{cc} + (L - M)\mu_a + M\mu_c.$$

Ground-state roots:  $\lambda = \mu_a = \mu_b = 0$




Figure : Ground-state root distribution for  $\lambda = 0$ ,  $N = 120$ .

# Large- $N$ limit

Ground-state root density:

$$\rho(v) = \sqrt{(b-v)(v-a)} \left( \frac{1}{2\pi M} + \frac{J+1}{2\pi M\sqrt{abv}} \right)$$

$$1 = \int_a^b dv \rho(v)$$

Ground-state energy:

$$E_0 = \mu \left( \frac{N+1}{2} + \frac{(J+1)^2}{2ab} \right) - \frac{(J+1)\mu}{2\alpha\sqrt{2N}\sqrt{ab}} \left( ab - \frac{(J+1)^2}{ab} \right)$$

Bethe ansatz equation:

$$\frac{J+1}{v} - v - \frac{\mu_c}{\Omega} = 2M \lim_{\epsilon \rightarrow 0} \int_a^{v-\epsilon} dw \frac{\rho(w)}{w-v} + 2M \lim_{\epsilon \rightarrow 0} \int_{v+\epsilon}^b dw \frac{\rho(w)}{w-v}$$

Links and Marquette, work in preparation

leads to

$$(\alpha b)^2 + 2(1 - \alpha^2)Nab - 4(J + 1)\alpha\sqrt{2N}\sqrt{ab} - 3(J + 1)^2 = 0.$$

For  $J = O(N^0)$ 

$$\alpha > 1 \quad \Rightarrow \quad ab \sim 2(\alpha^2 - 1)N,$$

$$\alpha = 1 \quad \Rightarrow \quad ab \sim 2^{5/3}(J + 1)^{2/3}N^{1/3},$$

$$\alpha < 1 \quad \Rightarrow \quad ab \sim (J + 1)^2 \left( \frac{2\sqrt{2}\alpha + \sqrt{2\alpha^2 + 6}}{2(1 - \alpha^2)} \right)^2 N^{-1}.$$

This yields

$$\alpha \geq 1 \quad \Rightarrow \quad \mathcal{O} = 1,$$

$$\alpha \leq 1 \quad \Rightarrow \quad \mathcal{O} = 1 + f^2 + \alpha \left( 2f + \frac{3}{\sqrt{2}\alpha} f^2 \right) \frac{df}{d\alpha}$$

where

$$f = \frac{2(1 - \alpha^2)}{2\sqrt{2}\alpha + \sqrt{2\alpha^2 + 6}}.$$


Molecular fraction:  $\lambda = \mu_a = \mu_b = 0$  - numerical

Figure : Order parameter. Inset shows the first derivative.

Links and Marquette, work in preparation

Molecular fraction:  $\lambda = \mu_a = \mu_b = 0$  - analytic curve for all finite  $J$  in the  $N \rightarrow \infty$  limit




Figure : Order parameter. Inset shows the first derivative.

# Future work

Extend the analysis to cover the entire parameter space.

